
Mitigating the risk of climate-induced slope instability from remote sensing multi-sensor data with artificial intelligence and numerical modelling

(Proposer: Filippo Catani)

Background and rationale

In the context of basin-scale modelling and assessment of slope instability, there is a relevant gap of knowledge which concerns the time component. Most of the field information relative to mass movements is only spatial and the great majority of the existing databases and inventories refer to a single event or are neglecting the time component (the so-called historical inventories), the main cause being a lack of a consistent and applicable methodology for collecting, checking, selecting, and normalizing the complex, multi-source information in a common standard including time of occurrence, geographic position and intensity estimation (Chae et al., 2017). In this framework, emerging technologies making use of machine intelligence are at the forefront in research innovation and provide a novel support to access, measure, and exploit time-related data sources previously not usable such as raw real-time sensor acquisitions, social networks content, continuous or discontinuous streaming from news and online databases. Among them, the automation of data mining of quantitative information from satellite, aerial and drone surveys, by using deep learning methods (DL) and recurrent neural networks (RNN) to perform classification, interpretation and forecasting of time series derived from monitoring of slope processes is of notable importance in the geosciences. Despite being very fashionable and promising, though, such technologies are still scantly used in engineering geology, with the exception of classical machine learning methods such as multi-variate regression or classification (Ermini et al., 2005; Catani et al., 2013), and an important research gap is yet present in the knowledge transfer from fields where DL and RNN are being routinely used, such as automotive and defense, to the realm of geomechanics and geotechnics applied to natural slopes. A few recent examples address the issue and put the basis for a new approach, by proposing novel concepts to model or forecast slope instability at various scales, going from satellite data interpretation (Mondini et al., 2011; Forzieri et al., 2012; Meena et al., 2023), to slope scale timeseries analysis (Turner et al., 2015; Nava et al., 2023), to the automated content extraction for the rapid alert and mapping of mass movements (Battistini et al., 2017; Catani, 2021; Nava et al., 2023, 2024).



Aim

The proposed PhD project builds on such recent advances related to the development of machine intelligence, to extend existing methods and techniques to the characterization of slope instability at the regional and basin scale for risk mitigation purposes with the help of multi-sensor data. Concisely, the main objectives of the PhD project will be: i) to exploit and adapt existing Deep Learning tools used for landslide detection from either optical or SAR data towards the integration on multi-sensor and fusion data for the development of advanced mapping techniques and towards data assimilation in numerical models for the spatial prediction of mass displacements, and, ii) to develop new methods based on Recurrent Neural Networks and time-series analysis for the forecasting of time-of-failure and the modelling and prediction of slope evolution in the short- to medium-term, both at the local scale and beyond, with a specific reference to mining-induced graound instability.

To pursue the objectives, the project foresees the continuation of new and promising research that applies DL to the automated recognition of mass movements in both optical and SAR imagery, including combinations of

them and data fusion with high-resolution topography, multi-temporal changes in the landscape morphometry and climate forcing.

Expected results

- i) Development of DL libraries and convolutional neural networks to extract quantitative information on active slope processes from combined Sentinel-1 + Sentinel-2 imagery;
- ii) Development of slope stability assessment classification tools starting from (i) with AI and numerical models;
- iii) Development of algorithms based on RNN for the semi-automated analysis of multi-point monitoring time-series to model displacement patterns and generate time forecasts.
- iv) Scaling-up methods and tools towards a global applicability of the most promising results.

Scientific cooperation

Most of the analysis and development of the PhD work will be carried out in cooperation with the CTTC (Centre Tecnològic de Telecomunicacions de Catalunya) (Prof. O. Monserrat) and the Cambridge University, Dept. of Earth Sciences (Prof. M. Van Wyk de Vries), partially under the ESA Geohazards Exploitation Platform (GEP, https://geohazards-tep.eu/#!) project for the development of downstream products of Sentinel constellations.

Funding and support

Funding will be provided by relevant connected projects such as the "ESA EGMS assessment program" sponsored by the Autorità di Distretto del Po, the "Return PNRR project", and the IPL-ISDR project "MUSE", sponsored by the International Consortium on Landslides.

References

- Battistini, A., Rosi, A., Segoni, S., Lagomarsino, D., Catani, F., Casagli, N., 2017. Validation of landslide hazard models using a semantic engine on online news. Applied Geography 82, 59–65. https://doi.org/10.1016/j.apgeog.2017.03.003 Catani, F., 2020. Landslide detection by deep learning of non-nadiral and crowdsourced optical images. Landslides. https://doi.org/10.1007/s10346-020-01513-4
- Catani, F., Lagomarsino, D., Segoni, S., Tofani, V., 2013. Landslide susceptibility estimation by random forests technique: sensitivity and scaling issues. Natural Hazards & Earth System Sciences 13.
- Chae, B.-G., Park, H.-J., Catani, F., Simoni, A., Berti, M., 2017. Landslide prediction, monitoring and early warning: a concise review of state-of-the-art. Geosciences Journal 21, 1033–1070. https://doi.org/10.1007/s12303-017-0034-4
- Ermini, L., Catani, F., Casagli, N., 2005. Artificial Neural Networks applied to landslide susceptibility assessment. Geomorphology 66(1–4), 327–343. https://doi.org/10.1016/j.geomorph.2004.09.025
- Forzieri, G., Moser, G., Catani, F., 2012. Assessment of hyperspectral MIVIS sensor capability for heterogeneous landscape classification. ISPRS Journal of Photogrammetry and Remote Sensing 74, 175–184. https://doi.org/10.1016/j.isprsjprs.2012.09.011
- Meena, S.R., Nava, L., Bhuyan, K., Puliero, S., Soares, L.P., Dias, H.C., Floris, M., Catani, F., 2023. HR-GLDD: a globally distributed dataset using generalized deep learning (DL) for rapid landslide mapping on high-resolution (HR) satellite imagery. Earth Syst. Sci. Data 15, 3283–3298. https://doi.org/10.5194/essd-15-3283-2023
- Mondini, A.C., Guzzetti, F., Reichenbach, P., Rossi, M., Cardinali, M., Ardizzone, F., 2011. Semi-automatic recognition and mapping of rainfall induced shallow landslides using optical satellite images. Remote Sensing of Environment 115, 1743–1757. https://doi.org/10.1016/j.rse.2011.03.006
- Nava, L., Monserrat, O., Catani, F., 2022. Improving Landslide Detection on SAR Data Through Deep Learning. IEEE Geosci. Remote Sensing Lett. 19, 1–5. https://doi.org/10.1109/LGRS.2021.3127073
- Nava, L., Carraro, E., Reyes-Carmona, C., Puliero, S., Bhuyan, K., Rosi, A., Monserrat, O., Floris, M., Meena, S.R., Galve, J.P., Catani, F., 2023. Landslide displacement forecasting using deep learning and monitoring data across selected sites. Landslides 20, 2111–2129. https://doi.org/10.1007/s10346-023-02104-9
- Nava, L., Tordesillas, A., Qian, G., Catani, F., 2024. Displacement residuals reveal landslide regime shifts. Landslides. https://doi.org/10.1007/s10346-024-02353-2
- Turner, D., Lucieer, A., de Jong, S., 2015. Time Series Analysis of Landslide Dynamics Using an Unmanned Aerial Vehicle (UAV). Remote Sensing 7, 1736–1757. https://doi.org/10.3390/rs70201736
- Yang, B., Yin, K., Lacasse, S., Liu, Z., 2019. Time series analysis and long short-term memory neural network to predict landslide displacement. Landslides 16, 677–694. https://doi.org/10.1007/s10346-018-01127-x